

- 353 Peng, S.C., Babcock, L.E., Cooper, R.A., 2012. The Cambrian Period. In Gradstein, F.M.,
354 Ogg, J.G., Schmitz, M.D., Ogg, G.M. eds., *The Geologic Time Scale 2012*, vol. 2.
355 Elsevier BV, Amsterdam, 437–488.
- 356 Pickett, J.W. and Jell, P.A., 1983. Middle Cambrian Sphinctozoa (Porifera) from New South
357 Wales. *Memoirs of the Association of Australasian Paleontologists*, 1: 85–92
- 358 Pouille, L., Obut O., Danelian T. et al., 2011, Lower Cambrian (Botomian) polycystine Radiolaria
359 from the Altai Mountains (southern Siberia, Russia). *Comptes Rendus Palevol*, 10(8): 627–633
- 360 Shu, D., Chen, L., 1989. Discovery of Early Cambrian radiolarian and its significance. *Science in
361 China*, 32(8): 986–994
- 362 White, R.D., 1986, Cambrian Radiolaria from Utah. *Journal of Paleontology*, 60(3): 778–780
- 363 Wrona, R. 2004. Cambrian microfossils from glacial erratics of King George Island, Antarctica.
364 *Acta Palaeontologica Polonica*, 49 (1): 13–56
- 365 Wu, W., Zhu, M., Steiner, M., 2014. Composition and tiering of the Cambrian sponge communities.
366 *Palaeogeography Palaeoclimatology Palaeoecology*, 398(398): 86–96
- 367 Won, M.Z., Below, R., 1999. Cambrian Radiolaria from the Georgina Basin, Queensland, Australia.
368 *Micropaleontology*, 45(4): 325–363
- 369 Yang, B., Steiner, M., Li, G., et al., 2014. Terreneuvian small shelly fauna of East Yunnan (South
370 China) and their biostratigraphic implications. *Palaeogeography Palaeoclimatology
371 Palaeoecology*, 398(2014): 28–58
- 372 Yin, H., Zeng, Y., Xia, W., 1994. Chert on the Southeast continental margin of the Yangtze platform.
373 *Acta Geologica Sinica*, 68(2): 132–141 (in Chinese with English abstract)
- 374 Yin, L., Wang C., Zhao, Y. et al., 2016. Early–Middle Cambrian Palynomorph Microfossils and
375 Related Geochemical Events in South China. *Journal of Earth Science*, 27(2): 180–186
- 376 Zhang, L., Danelian, T., Feng, Q.L., et al., 2003. On the Lower Cambrian biotic and geochemical
377 record of the Hetang Formation (Yangtze Platform, South China): evidence for biogenic silica
378 and possible presence of Radiolaria. *Journal of Micropaleontology*, 32(2): 207–217
- 379 Zhang, X., Aldridge, R. J., 2007. Development and diversification of trunk plates of the lower
380 Cambrian lobopodians. *Paleontology*, 50(2): 401–415
- 381 Zhang, X., Cui, L., 2016. Oxygen requirements for the Cambrian explosion. *Journal of Earth
382 Science*, 27(2): 187–195
- 383 Zhang, X., Pratt, B., 1994. Rare and extraordinary Early Cambrian sponge spicule assemblage from
384 China. *Geology*, 22(1): 43–46
- 385 Zhang, X., Shu, D., Han, J., et al., 2014. Triggers for the Cambrian explosion: Hypotheses and
386 problems. *Gondwana Research*, 25(3): 896–909
- 387 Zhao, G., 1999. The influence of biogenic procession on the accumulation and precipitation of
388 silica—evidence from south of Anhui and west of Zhejiang. *Acta Sedimentologica Sinica*,
389 17(1): 30–37 (in Chinese with English abstract)
- 390 Zheng, N., Song, T., Li, Y., et al., 2012. The discovery of the Lower Cambrian and Middle
391 Ordovician Radiolaria in the South China orogenic belt. *Geology in China*, 39(1): 260–265 (in
392 Chinese with English abstract)
- 393 Zhou C., Jiang S., 2009. Palaeoceanographic redox environments for the lower Cambrian Hetang
394 Formation in South China: Evidence from pyrite framboids, redox sensitive trace elements, and
395 sponge biota occurrence. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 271(4): 279–
396 286