

- 353 Peng, S.C., Babcock, L.E., Cooper, R.A., 2012. The Cambrian Period. In Gradstein, F.M.,
354 Ogg, J.G., Schmitz, M.D., Ogg, G.M. eds., The Geologic Time Scale 2012, vol. 2.
355 Elsevier BV, Amsterdam, 437–488.
- 356 Pickett, J.W. and Jell, P.A., 1983. Middle Cambrian Sphinctozoa (Porifera) from New South
357 Wales. *Memoirs of the Association of Australasian Paleontologists*, 1: 85–92
- 358 Pouille, L., Obut O., Danelian T. et al., 2011, Lower Cambrian (Botomian) polycystine Radiolaria
359 from the Altai Mountains (southern Siberia, Russia). *Comptes Rendus Palevol*, 10(8): 627–633
- 360 Shu, D., Chen, L., 1989. Discovery of Early Cambrian radiolarian and its significance. *Science in
361 China*, 32(8): 986–994
- 362 White, R.D., 1986, Cambrian Radiolaria from Utah. *Journal of Paleontology*, 60(3): 778–780
- 363 Wrona, R. 2004. Cambrian microfossils from glacial erratics of King George Island, Antarctica.
364 *Acta Palaeontologica Polonica*, 49 (1): 13–56
- 365 Wu, W., Zhu, M., Steiner, M., 2014. Composition and tiering of the Cambrian sponge communities.
366 *Palaeogeography Palaeoclimatology Palaeoecology*, 398(398): 86–96
- 367 Won, M.Z., Below, R., 1999. Cambrian Radiolaria from the Georgina Basin, Queensland, Australia.
368 *Micropaleontology*, 45(4): 325–363
- 369 Yang, B., Steiner, M., Li, G., et al., 2014. Terreneuvian small shelly fauna of East Yunnan (South
370 China) and their biostratigraphic implications. *Palaeogeography Palaeoclimatology
371 Palaeoecology*, 398(2014): 28–58
- 372 Yin, H., Zeng, Y., Xia, W., 1994. Chert on the Southeast continental margin of the Yangtze platform.
373 *Acta Geologica Sinica*, 68(2): 132–141 (in Chinese with English abstract)
- 374 Yin, L., Wang C., Zhao, Y. et al., 2016. Early–Middle Cambrian Palynomorph Microfossils and
375 Related Geochemical Events in South China. *Journal of Earth Science*, 27(2): 180–186
- 376 Zhang, L., Danelian, T., Feng, Q.L., et al., 2003. On the Lower Cambrian biotic and geochemical
377 record of the Hetang Formation (Yangtze Platform, South China): evidence for biogenic silica
378 and possible presence of Radiolaria. *Journal of Micropaleontology*, 32(2): 207–217
- 379 Zhang, X., Aldridge, R. J., 2007. Development and diversification of trunk plates of the lower
380 Cambrian lobopodians. *Paleontology*, 50(2): 401–415
- 381 Zhang, X., Cui, L., 2016. Oxygen requirements for the Cambrian explosion. *Journal of Earth
382 Science*, 27(2): 187–195
- 383 Zhang, X., Pratt, B., 1994. Early and extraordinary Early Cambrian sponge spicule assemblage from
384 China. *Geology*, 22(1): 43–46
- 385 Zhang, X., Shu, D., Han, J., et al., 2014. Triggers for the Cambrian explosion: Hypotheses and
386 problems. *Gondwana Research*, 25(3): 896–909
- 387 Zhao, G., 1999. The influence of biogenic procession on the accumulation and precipitation of
388 silica—evidence from south of Anhui and west of Zhejiang. *Acta Sedimentologica Sinica*,
389 17(1): 30–37 (in Chinese with English abstract)
- 390 Zheng, N., Song, T., Li, Y., et al., 2012. The discovery of the Lower Cambrian and Middle
391 Ordovician Radiolaria in the South China orogenic belt. *Geology in China*, 39(1): 260–265 (in
392 Chinese with English abstract)
- 393 Zhou C., Jiang S., 2009. Palaeoceanographic redox environments for the lower Cambrian Hetang
394 Formation in South China: Evidence from pyrite framboids, redox sensitive trace elements, and
395 sponge biota occurrence. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 271(4): 279–
396 286