

- Mügler, I., Gleixner, G., Günther, F., et al., 2010. A multi-proxy approach to reconstruct hydrological changes and Holocene climate development of Nam Co, Central Tibet. *Journal of Paleolimnology*, 43 (4): 625–648
- Murray, A.S., Wintle, A.G., 2000. Luminescence dating of quartz using an improved single-aliquot regenerative-dose protocol. *Radiation Measurement*, 32 (1): 53–57
- Murray, A.S., Wintle, A.G., 2003. The single aliquot regenerative dose protocol: potential for improvements in reliability. *Radiation Measurements*, 37 (37): 377–381
- Prescott, J., Hutton, J.T., 1994. Cosmic ray contributions to dose rates for luminescence and ESR dating: large depths and long-term time variations. *Radiation Measurements*, 23 (2): 497–500
- Ramisch, A., Lockot, G., Haberzettl, T., et al., 2016. A persistent northern boundary of Indian Summer Monsoon precipitation over Central Asia during the Holocene. *Scientific Reports*, 6: 25791, DOI: 10.1038/srep25791
- Ran, M., Feng, Z., 2013. Holocene moisture variations across China and driving mechanisms: A synthesis of climatic records. *Quaternary International*, 313–314: 179–193
- Reimer, P.J., Baillie, M.G.L., Bard, E., et al., 2004. IntCal04 terrestrial radiocarbon age calibration, 0–26 cal kyr BP. *Radiocarbon*, 46 (3): 1029–1058
- Shen, Z.X., Bloemendal, J., Mauz, B., et al., 2008. Holocene environmental reconstruction of sediment-source linkages at Crummock Water, English Lake District, based on magnetic measurements. *The Holocene*, 18 (18): 129–140
- Stauch, G., Schulte, P., Ramisch, A., Hartmann, K., Hülle, D., Lockot, G., Diekmann, B., Nottebaum, V., Müller, C., Wünnemann, B., Yan, D., Lehmkuhl, F., 2017. Landscape and climate on the northern Tibetan Plateau during the late Quaternary. *Geomorphology*, 286: 78–92
- Thomas, P.J., Murray, A.S., Sandgren, P., 2003. Age limit and age underestimation using different OSL signals from lacustrine quartz and polymineral fine grains. *Quaternary Science Review*, 22 (10): 1139–1143
- Tian, L., Yao, T., MacClune, K., et al., 2007. Stable isotopic variations in west China: A consideration of moisture sources. *Journal of Geophysical Research*, 112 (D10): 185–194
- Wang, A., Smith, J.A., Wang, G., et al., 2009. Late Quaternary river terrace sequences in the eastern Kunlun Range northern Tibet: A combined record of climatic change and surface uplift. *Journal of Asian Earth Sciences*, 34 (4): 532–543
- Wang, W., Feng, Z., Ran, M., et al., 2013. Holocene climate and vegetation changes inferred from pollen records of Lake Aibi, northern Xinjiang, China: A potential contribution to understanding of Holocene climate pattern in East-central Asia. *Quaternary International*, 311: 54–62
- Wang, X., Zhou, A., Sun, Z., 2016. Spatial and temporal dynamics of lakes in Nam Co Basin, 1991–2011. *Journal of Earth Science*, 27 (1): 130–138
- Wang, Y., Shen, J., Wu, J.L., et al., 2007. Hard-water effect correction of lacustrine sediment ages using the relationship between ^{14}C levels in lake waters and in the atmosphere: The case of Lake Qinghai. *Journal of Lake Sciences*, 19 (5): 504–508 (in Chinese with English Abstract)
- Watanabe, T., Matsunaka, T., Nakamura, T., et al., 2010. Changes of organic matter sources in sediment core from a high-altitude lake (Pumoyum Co, southeastern Tibetan Plateau) over the last 19000 years. *Nuclear Instruments and Methods in Physics Research*, 268 (7–8): 1070–1072
- Wintle, A.G., Murray, A.S., 2006. A review of quartz optically stimulated luminescence characteristics and their relevance in single-aliquot regeneration dating protocols. *Radiation*

Measurements, 41, 369–391

- Wu, Y., Wang, S., Zhou, L., 2011. Possible factors causing older radiocarbon age for bulk organic matter in sediment from Daihai Lake, north China. *Radiocarbon*, 53 (2): 359–366
- Yang, B., Qin, C., Wang, J.L., et al., 2014. A 3,500-year tree-ring record of annual precipitation on the northeastern Tibetan Plateau. *Proceedings of the National Academy of Sciences of the United States of America*, 111 (8): 2903–2908
- Yang, K., Wu, H., Qin, J., et al., 2014. Recent climate changes over the Tibetan Plateau and their impacts on energy and water cycle: A review. *Global and Planetary Change*, 112 (1): 79–91
- Yan, D., Wünnemann, B., 2014. Late Quaternary water depth changes in Hala Lake, northeastern Tibetan Plateau, derived from ostracod assemblages and sediment properties in multiple sediment records. *Quaternary Science Review*, 95 (95): 95–114
- Yu, L.P., Lai, Z.P., 2012. OSL chronology and palaeoclimatic implication of aeolian sediments in the eastern Qaidam Basin of the northeastern Qinghai-Tibetan Plateau. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 337–338 (4): 120–129
- Yu, L.P., Lai, Z.P., 2014. Holocene climate change inferred from stratigraphy and OSL chronology of aeolian sediments in the Qaidam Basin, northeastern Qinghai-Tibetan Plateau. *Quaternary Research*, 81 (3): 488–499
- Zhang, J., Chen, F., Holmes, J.A., et al., 2011. Holocene monsoon climate documented by oxygen and carbon isotopes from lake sediments and peat bogs in China: a review and synthesis. *Quaternary Science Reviews*, 30 (15–16): 1973–1987
- Zhang, W., Mischke, S., Zhang, C., et al., 2013. Ostracod distribution and habitat relationships in the Kunlun Mountains, northern Tibetan Plateau. *Quaternary International*, 313–314 (10): 38–46
- Zhao, Y., Yu, Z., Chen, F., et al., 2007. Holocene vegetation and climate history at Hurleg Lake in the Qaidam Basin, northwest China. *Review of Palaeobotany and Palynology*, 145 (3–4): 275–288
- Zhou, A., Chen, F., Wang Z., et al., 2009. Temporal change of radiocarbon reservoir effect in Suga Lake northwest China during the Lake Holocene. *Radiocarbon*, 51 (2): 529–535
- Zhu, L.P., Wu, Y.H., Wang, J.B., et al., 2008. Environmental changes since 8.4 ka reflected in the lacustrine core sediments from Nam Co, central Tibetan Plateau. *The Holocene*, 18 (5): 831–839
- Zeng, F., Xiang, S., 2015. Geochronology and mineral composition of the Pleistocene sediments in Xitaijinair salt lake region. *Journal of Earth Science*, 28 (4): 1–6
- Zeng, F., Liang, M., Peng, S., et al., 2015. Sr-Nd-Pb Isotopic Compositions of the Neogene Eolian Deposits in the Xining Basin and Implications for Their Dust Sources. *Journal of Earth Science*, 26 (5): 669–676